





Digital Version



You will need to shower with a special soap called chlorhexidine gluconate. The purpose of the chlorhexidine gluconate soap is to reduce the normal bacteria on your skin, that may be a potential source of infection thus it reduces chances of post-surgery infection. The soap may come in a liquid form.

WHEN TO HAVE BATH?

Whole body wash should be carried out 2 times:

- Previous day of the surgery in the night.
- In the morning on the day of surgery (Post clipping of hair)

HOW TO DO IT?

- Please remove all body piercing jewelries prior to shower and remain ewelry free until after surgery.
- Take normal head to toe bath with regular soap and water before chlorhexidine bath.
- Apply 40-50 ml of undiluted chlorhexidine soap to your entire body from head to toe.
- Pay special attention to the area of surgery and also armpit and groin.
- Turn water off to allow lather to remain in the body for 5 minutes, then wash your body thoroughly.
- After final rinse, pat yourself dry with a clean towel and put on clean cloths.

DON'T

- Do not use chlorhexidine if you are allergic to it.
- Do not use chlorhexidine on your face, private part and any cut onyour skin.
- Do not scrub your skin too hard after applying chlorhexidine.
- Do not use regular soap after chlorhexidine is used.
- Do not apply lotion, powder, after shave, deodorant, perfume on the skin after bathing with chlorhexidine.





PLEASE SCAN FOR THE LOCATION



Mankhool, Kuwait Road, Al Mankhool - Dubai © 04 440 0500 ⊕ asterhospitals.com ④ ⑨ ⊚ asterhospital

DEPARTMENTS

Accident & Emergency · Anesthesiology · Cardiology · Clinical Nutrition

Dental Centre · Dermatology · ENT · General Surgery · Insurance

Internal Medicine · Laboratory Services · Neurosurgery · Ophthalmology

Obstetrics & Gynaecology · Orthopedics · Patient Affairs