



UREA BREATH TEST







Digital Version



• FAST for at least one (1) hour before the test. This means that you consume no food and drink only water prior to the test.

• If you have Phenylketonuric should avoid the test. Active ingredient is mixed with aspartame and contains 84 mg phenylalanine per dosage unit.

• Avoid proton pump inhibitors including Prilosec, Prevacid, Nexium, Protonix, Aciphex, and Dexilant within two (2) weeks prior to the urea breath test.

• Avoid antibiotics two (2) weeks prior to the urea breath test.

• Avoid bismuth preparations (Pepto Bismol) within two (2) weeks prior to the urea breath test.

• Do not test before four (4) weeks after H. pylori treatment.

• Prior to testing, the patient may take H2 antagonists (Tagamet, Zantac, Pepcid, Axid) or Antacids (Tums, Rolaids, Maalox).

PRECAUTIONS

• Not re-capping the breath collection bag properly. The white cap must snap in place.

• Administration of Pranactin-Citric solution before collecting the baseline breath sample.

• Consumption of food, beverages, or smoking during the 15-minute breath collection interval.

POTENTIAL COMPLICATIONS

• During post-approval use of the urea breath test, the following adverse events have been identified: anaphylactic reaction, hypersensitivity, rash, burning sensation in the stomach, tingling in the skin, vomiting and diarrhea. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to establish a causal relationship to drug exposure.





PLEASE SCAN FOR THE LOCATION



Mankhool, Kuwait Road, Al Mankhool - Dubai © 04 440 0500 ⊕ asterhospitals.com ④ ⑨ ◎ asterhospital

DEPARTMENTS

Accident & Emergency · Anesthesiology · Cardiology · Clinical Nutrition Dental Centre · Dermatology · ENT · General Surgery · Insurance Internal Medicine · Laboratory Services · Neurosurgery · Ophthalmology Obstetrics & Gynaecology · Orthopedics · Patient Affairs